

Environmental Fact Sheet (# 33)

Crude Palm Kernel Oil (CPKO)

renewable precursor

Substance Identification	
IUPAC Name	Oils, palm kernel
CAS Number	8023-79-8
Other Names	
Molecular Formula	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>UVCB</p> </div> <div style="text-align: center;"> $\begin{array}{c} \text{O} \\ \parallel \\ \text{CH}_2 - \text{O} - \text{C} - \text{R} \\ \\ \text{CH}_2 - \text{O} - \text{C} - \text{R}' \\ \\ \text{O} \\ \parallel \\ \text{CH}_2 - \text{O} - \text{C} - \text{R}'' \end{array}$ </div> </div>
Physical/Chemical Properties [1]	
Fatty acid composition	85 % saturated Fatty acid (Lauric (40-52), Myristic (14-18), Palmitic (7-9), Capric (3-7), Caprylic (3-5) and others in lower composition) and 15% unsaturated fatty acid (Oleic (11-19), Palmitoleic (0-1), Linoleic (0.5-2))
Molecular Weight	No data available
Physical state	No data available
Appearance	No data available
Odour	No data available
Density	0.893 at 60 °C
Melting Points	22-26 °C
Boiling point	No data available
Flash Point	No data available
Vapour Pressure	No data available
Water Solubility	No data available
Viscosity	30.92 cSt at 37.8°C
Flammability	No data available
Explosive Properties	No data available
Surface Tension	No data available
Octanol/water Partition coefficient (Kow)	No data available
Product and Process Description	<p>The oil palm (<i>Elaeis guineensis</i>) is cultivated mainly for its fruit which is a source of vegetable oil and fat. The oil palm fruits (stone fruit) are harvested in fresh fruit bunches. The processing of palm kernel oil contains different steps:</p> <ol style="list-style-type: none"> 1) Reception of FFBS from the palm oil plantations. 2) Sterilisation, where vapour helps to loosen the individual fruits from the bunch and also deactivates the enzymes which cause the breakdown of the oil into free fatty acids.

	3) Stripping of bunches: the fruitlets are separated from bunches which are then called “empty fruit bunches. 4) Digestion: the empty fruit bunches are converted into a homogenous oily mash by a digester. 5) Kernel extraction: in the oil extraction of the palm oil, the kernel is obtained as a co-product. 6) Cleaning of the kernel 7) Palm kernel oil extraction: the palm kernel oil is extracted from the kernels using a screw press. 8) Purification: oil is purified through a coarse screen filter and a filter press. Palm kernel cake is obtained as by-product, which can be used as animal feed.
Application	Palm kernel oil derivatives are mostly used as soap production, margarine, lubricating oil production, insecticides production, vegetable oil-based printing inks production, food and cosmetic industry and other applications [4][5].

Life Cycle Assessment

General Introduction

These Environmental Fact Sheets are a product of the *ERASM Surfactant Life Cycle & Ecofootprinting (SLE)* project. The objective of this project was to create representative, global, attributional, industry average datasets for oil palm-based precursors in the production of several particular surfactants.

The Fact Sheets are based upon life cycle assessment (LCA) and have been prepared in accordance with the ISO standard [ISO 14040: 2006 and ISO 14044: 2006]. In addition, the project follows the ILCD (2010) handbook. This Fact Sheet describes the cradle-to-gate production for Crude palm kernel Oil (CPKO). CPKO is an oleo chemical precursor for the production of surfactants.

Further information on the ERASM SLE project and the source of these datasets can be found in [2].

The full LCI can be accessed via www.erasm.org or via <http://lcdn.thinkstep.com/Node/>

Goal and Scope of ERASM SLE Project [2]

The main goal was to develop the LCI datasets for the production of crude palm kernel oil (CPKO) for further use in LCA studies.

Temporal Coverage	Data collected for the production were based on literature research covering one production year technology. Due to lack of data, a consistent usage of data with one reference year or reference time was not possible. Some parameters used do not change over time. All used values have been published 2007 or after. Time dependent inputs, like yield are averaged over several years to be able to exclude peaks or years with poor harvests. Background data dates back to the reference year 2011. The datasets are considered to be valid until substantial technological changes in the production chain occur. The temporal representativeness for CPKO was considered ‘good’
Geographical Coverage	Data for palm kernel oil came from the two most important global suppliers of palm kernel oil, Indonesia and Malaysia, covering approximately 83% of the global production. In the background system some processes are modeled for different geographical system boundaries (mostly Germany and the US). The geographical representativeness for CPKO was considered ‘fair’.
Technological Coverage	The production technology was based on different literature sources. Partly processes with better data availability were chosen to be used as representative technology rather than creating averages of different technologies. I.e. for palm kernel oil mills only the technology of mechanical extraction was applied and the location of the palm kernel oil mill was set to be close to the sea and not located next to the palm oil mill. In the palm kernel oil refinery only physical processing was considered. As the palm oil industry is rapidly expanding and evolving, full technological representativeness cannot be guaranteed. All relevant technologies were considered and technology modelled to best knowledge. The technological representativeness for CPKO was considered ‘good’. Figure 1 provides a schematic overview of the production process of CPKO.

Declared Unit	In ERASM SLE project the declared unit (functional unit) and the reference flow of the product is one thousand kilogram (1000 kg). This was the reference unit also used in [3]. Functional Unit: 1 tonne 'Crude Palm Kernel Oil', at the palm oil kernel mill gate															
Cradle-to Gate System Boundaries	<table border="1"> <thead> <tr> <th style="background-color: #ADD8E6;">Included</th> <th style="background-color: #ADD8E6;">Excluded</th> </tr> </thead> <tbody> <tr> <td>Palm Oil Plantation</td> <td>Palm oil nursery</td> </tr> <tr> <td>Direct Land use change effects</td> <td>Equipment</td> </tr> <tr> <td>Unit processes (including material and energy inputs) of the palm oil mill</td> <td>Transport to Europe</td> </tr> <tr> <td>Unit processes (including material and energy inputs) of the palm kernel oil mill</td> <td></td> </tr> <tr> <td>Transport from the field to the palm oil mill</td> <td></td> </tr> <tr> <td>Transport from the palm oil mill to the palm kernel oil mill</td> <td></td> </tr> </tbody> </table>		Included	Excluded	Palm Oil Plantation	Palm oil nursery	Direct Land use change effects	Equipment	Unit processes (including material and energy inputs) of the palm oil mill	Transport to Europe	Unit processes (including material and energy inputs) of the palm kernel oil mill		Transport from the field to the palm oil mill		Transport from the palm oil mill to the palm kernel oil mill	
	Included	Excluded														
	Palm Oil Plantation	Palm oil nursery														
	Direct Land use change effects	Equipment														
	Unit processes (including material and energy inputs) of the palm oil mill	Transport to Europe														
	Unit processes (including material and energy inputs) of the palm kernel oil mill															
	Transport from the field to the palm oil mill															
Transport from the palm oil mill to the palm kernel oil mill																
Assumptions and Limitations	<p>The major assumptions made on land use change cultivation, processing usage of co-products, emissions from land use change and cultivation on peatland.</p> <p>Exclusion of primary data collection as it would not be feasible to assess primary data of the entire global production of palm kernel oil products. There is a lack of information, so several flows were only estimated (i.e. amount waste water or the exact composition of POME). The literature sources used could not display all technologies and production pathways. Model agrarian systems are considered but may strongly vary within one region. A mix of Malaysian and Indonesian palm kernel oil was used to describe a global average of these products (For Indonesia there were big data gaps so that the same process conditions as for Malaysia are assumed).</p>															
Cut-off Criteria	<p>The LCI study included all the flows which had more than 2% of the cumulative mass inputs and greater than 2 % of the cumulative energy of the respective gate-to-gate model inventory, providing its environmental relevance is not a concern. Material flows which leave the system (emissions) and whose environmental impact was greater than 2% of the whole impact of an impact category that has considered in the assessment were covered. The sum of the neglected material flows must not exceed 5% of mass, energy or environmental relevance of the system inventory.</p>															
Calculation Rules	Allocation	Mass allocation was applied for palm kernel oil between the product crude palm kernel oil and the co-product palm kernel cake.														
	Aggregated data	From literature research.														
Life Cycle Inventory and Impact Assessment [2]																
<p>Based on the LCI data an environmental impact assessment was performed for the indicators Primary Energy Demand (PED), Global Warming Potential (GWP), Eutrophication Potential (EP) and Acidification Potential (AP). Other impacts may be calculated from the full LCI dataset. Table 1 shows the environmental impact results per 1 tonne of CPKO.</p> <p><u>Primary Energy Demand (PED)</u>: An analysis of the inventory data showed that the PED come from electricity which is required in the palm oil mill, palm kernel oil mill.</p> <p><u>Global Warming Potential (GWP)</u>: An analysis of the inventory data shows that the main contribution comes from cultivation on peat land. The second biggest contributor of carbon dioxide emissions are the ones which occur due to land use change. Cultivation and processing contributes most methane emissions. Methane emissions occur to approximately 82-84% in the palm oil mill and originate from the POME treatment. Other methane emissions occur on the plantation. Nitrous oxide emissions occur almost entirely on the plantation.</p> <p><u>Air emissions related to Acidification (AP) and Eutrophication Potential (EP)</u>: An analysis of the inventory data shows that ammonia, nitrogen oxides, and sulphur dioxide emissions from LUC originate from the incineration of vegetation of the above ground biomass and of shells and fibres at the palm oil mill.</p> <p>Besides ammonia emissions occur during cultivation and POME treatment. Hydrogen sulphide is emitted during the provision of electricity which is required in the palm oil mill and palm kernel oil mill. Nitrogen monoxides are emitted during the transport of intermediate products and in the plantation (originate from agricultural processes). Nitrogen oxide emissions occur during the provision of electricity, the combustion of shells in the palm oil mill and use of tractors on the field.</p> <p><u>Emissions to fresh water related to Acidification (AP) and Eutrophication Potential (EP)</u>: An analysis of the inventory data</p>																

showed that the main emissions (nitrogen and phosphorus) to the fresh water occur during the cultivation of palm oil.

Sensitivity Analysis: Different sensitivity analysis were carried out. The analyses showed that allocation method, yield, fertilizer use, land use change have an impact on the results. Particularly in the case of economic allocation, the choice is strongly affecting the overall results.

Table 1. Global warming potential, acidification potential and eutrophication potential related to emissions to air and fresh water, and primary energy demand per 1 tonne of CPKO

LCI result	Unit	Amount		
		Cultivation and processing	Land Use Change	Cultivation on peatland
Air emissions related to Global Warming Potential				
Carbon uptake, biotic	kg	-5.04E+03	-	-
Carbon dioxide, fossil	kg	3.06E+02	1.05E+03	2.74E+03
Carbon dioxide, biotic	kg	2.37E+03	-	-
Methane	kg	3.48E+01	-	-
Nitrous oxide (laughing gas)	kg	2.30E-01	-	7.81E-01
Air emissions related to Acidification and Eutrophication Potential				
Ammonia	kg	1.07E-01	7.27E-01	-
Hydrogen sulfide	kg	2.21E-01	-	-
Nitrogen monoxide	kg	2.78E-01	-	-
Nitrogen oxides	kg	4.88E+00	1.76E+00	-
Sulphur dioxide	kg	1.55E+00	8.39E-01	-
Emissions to fresh water related to Eutrophication Potential				
Nitrogen organic bound	kg	4.92E-01	-	-
Phosphate	kg	1.18E-01	-	-
Total GWP (according to [IPCC 2007])	kg	-9.52E+02	1.05E+03	2.97E+03
<i>EP</i>	kg	1.27E+00	-	2.11E-01
<i>AP</i>	kg	6.61E+00	-	-
Primary energy demand				
Primary energy demand from renewable materials (net calorific value)	MJ		4.92E+04	
Primary energy demand from fossil materials (net calorific value)	MJ		5.01E+03	
Primary energy demand from fossil and renewable materials (net calorific value)	MJ		5.42E+04	

References for the ERASM SLE Project	
Data Owner and Commissioner of the study	ERASM (Environment & Health Risk Assessment and Management). A research partnership of the Detergents and Surfactants Industries in Europe (www.erasm.org).
LCA Practitioner	thinkstep AG (www.thinkstep.com)
Reviewers	Prof. Walter Kloepffer, LCA Consult Mrs. Charlotte Petiot and Dr. Yannick Leguern, BioS by Deloitte Jannick Schmidt (2.0 LCA Consultants)
References	<p>[1] Daniel Swern (1979). Bailey's Industrial oil fat products, volume 1, 1979.</p> <p>[2] Schowanek. D <i>et al.</i> (2017). New and Updated Life Cycle Inventories for Surfactants used in European Detergents: Summary of the ERASM Surfactant Life Cycle and Ecofootprinting Project. Int J. LCA, in press.</p> <p>[3] CEFIC-Franklin (1994). Resource and environmental profile analysis of petrochemical and oleo chemical surfactants produced in Europe. Phase II Final Report, Franklin Associates, LTD.</p> <p>[4] Ajiwe <i>et al.</i> (2003). Biodiesel Fuels from Palm Oil, Palm Oil Methylene Ester and Ester-Diesel Blends, Bull. Chem Soc. Ethiop. 17, 19-26.</p> <p>[5] Ghazali <i>et al.</i> (2005). Non-Food Applications of Palm-Based Products – Market Opportunities and Environmental Benefits. Palm Oil Developments 44.</p> <p>[6] Subramanian <i>et al.</i> (2010). Life Cycle Assessment of Production of Crude Palm Kernel Oil (Part 3a). J. Oil pal Research, 22, 895-903.</p>

Figure 1. Production process of Palm Kernel Oil [6].

